



Observational Study

Vascular risk factors and carotid atheromatous disease in patients over 65 years of age

Sónia Mateus^{1,2*}, Vanda Pós de Mina¹, Patrícia Coelho²⁻⁴, Francisco Rodrigues²⁻⁴ and Irene Mendes¹

¹Cardiac and Neurovascular Ultrasound Laboratory, Hospital Espírito Santo de Évora, Évora, Portugal

²Castelo Branco Polytechnic Institute, Higher School of Health Dr. Lopes Dias, Castelo Branco, Portugal

³Integrated Member of the Sport Research Unit, Health S. Exercise Unit (SHERU), Portugal

⁴Integrated Member Research Unit Quality of Life in the Rural World (Qrural), Portugal

Received: 10 August, 2023

Accepted: 22 August, 2023


Published: 23 August, 2023

*Corresponding author: Sónia Mateus, Ultrasound Technician, Cardiac and Neurovascular Ultrasound Laboratory, Hospital Espírito Santo de Évora, Évora, Portugal, Castelo Branco Polytechnic Institute, Higher School of Health Dr. Lopes Dias, Castelo Branco, Portugal, Tel: 932334863; E-mail: soniamatildemateus@gmail.com

Keywords: Carotid atherosclerosis; Cardiovascular risk factors; Carotid intima-media thickness

Copyright License: © 2023 Mateus S, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

<https://www.peertechzpublications.org>



Abstract

Introduction: Vascular risk factors are decisive in the evolution of atherosclerotic disease and the carotid and vertebral Doppler ultrasound allows monitoring its onset and progress. The measurement of arterial wall thickening allows the early diagnosis of the disease enhancing its treatment and control of vascular risk factors.

Aim: Analysing the presence of atheromatous disease in individuals aged > 65 years and understanding its correlation with vascular risk factors.

Materials and methods: This is a cross-sectional observational study, in individuals aged > 65 years and underwent carotid echoDoppler between January 1, 2012, and December 31, 2021. The intimal-media index was calculated, as was the presence of atheromatous plaques, their hemodynamic repercussion and vascular risk factors were recorded.

Results: A sample of 5885 individuals was obtained with 41.8% female and 58.2% male. The mean age was 76.59 (\pm 6.69), with a range between 65 and 98 years. Arterial hypertension was the most prevalent 81.3%. There was a significant positive relationship between the intima-media index and age ($p = 0.001$). In the presence of plaques, male gender, arterial hypertension, diabetes, dyslipidemia, and smoking it presents a positive correlation.

Conclusion: In this sample, non-modifiable vascular risk factors seem to be determinants in the presence of increased arterial wall thickness. In the presence of signs of a more advanced stage of atherosclerotic disease, modifiable vascular risk factors are decisive, corroborating the already-known importance of strict control over them for their treatment.

Introduction

Vascular risk factors that contribute to atherosclerotic disease can be non-modifiable such as age, ethnicity, gender, and family history of atherosclerosis, or modifiable risk factors, which can be controlled through re-education for healthy lifestyles, such as arterial hypertension, Diabetes Mellitus (DM), hypercholesterolemia, hypertriglyceridemia, obesity, smoking, sedentary lifestyle, excessive alcohol consumption and embolic heart diseases (such as Atrial Fibrillation (AF) or patent foramen ovale) [1-5].

Atherosclerosis is an inflammation that occurs mainly in the arteries of medium and large calibre, in a chronic and progressive way, due to the progressive and/or sudden change in the composition of the blood, through the effect of risk factors, resulting in damage to the vessel wall. This inflammation will result in the increasing deposit of lipids, calcium, blood cells, and constituents in the vessel walls, forming atheromatous plaques that, in more severe cases, affect blood circulation and lead to ischemia in the dependent territory [6,7].

Atherosclerosis is the etiology that can affect the coronary

arteries, with a risk of angina pectoris and/or acute myocardial infarction, it also affects the arteries of the limbs resulting in peripheral vascular disease and affects extra and/or intracranial arteries, being a factor risk of ischemic stroke [6,7]. The highest occurrence of this pathology type, as a rule, occurs in the elderly population over 65 years and in males [4].

Carotid arteries are among the vessels most affected by atherosclerotic disease due to their size and turbulence of flow at the bifurcation. That is why it is extremely important to have an early perception of the presence of atheromatous disease, which is characterized in an initial phase by lipid infiltration, later evolving into atheromatous plaques and reduction of the carotid lumen due to stenosis. This study is performed by cervical Doppler ultrasound of the carotid and vertebral arteries, allowing the quantification of arterial wall thickening called the Intima-Media Index (IMI), which has great potential for the non-invasive assessment of atherosclerotic disease, as well as the presence of atheromatous disease in more advanced stages, already with the presence of atheromatous plaques and respective hemodynamic repercussions. Several studies point to a relationship between the incidence of cardio and cerebrovascular diseases and increased IMI values. This represents the earliest form of atheromatous disease, being a risk marker for Acute Myocardial Infarction (AMI) and stroke, which justifies continued research and interest in this topic [8-10]. Cardio and cerebrovascular diseases of atheromatous origin are the main cause of mortality and morbidity in Portugal, namely in the Alentejo region. This is a particular region because of its own gastronomic and demographic characteristics, namely with high-fat levels and a very aged population [3,11,12]. We consider the over 65 years old as the elderly cut-off in Portugal [11]. With that premise, we designed this study aiming to analyse the presence of atheromatous disease in a group of individuals aged more than 65 years old from the Alentejo region and understand its correlation with vascular risk factors.

Materials and methods

Study design

The study was performed with a non-probabilistic sample for convenience consisting of 5885 individuals aged 65 years or older who underwent the cervical Doppler ultrasound examination in the central hospital at Alentejo – Portugal, between January 2002 and December 2021, and about which it was possible to collect information on cardiovascular risk factors, as well as Doppler ultrasound findings on the carotid and vertebral arteries and corresponding IMI values.

Location

Ultrasound lab in central hospital at Alentejo – Portugal.

Sample population and size

The sample consisted of 5885 individuals.

Selection criteria

All the individuals selected were aged 65 years or older and underwent the cervical Doppler ultrasound examination in the

central hospital at Alentejo – Portugal between December 2021 and January 2002.

Data collection

The variables collected for the study were gender, age, race, family history of AMI and stroke, presence of hypertension, smoking habits, presence of dyslipidemia, AF, presence of other embolic heart diseases, presence of diabetes mellitus, IMI and Doppler ultrasound results of the carotid axes (right and left) and vertebral arteries. Qualitative variables (arterial hypertension, dyslipidemia, AF, DM, and other embolic heart diseases) were subdivided into “yes” which corresponds to the presence of co-morbidity in the individual and “no” to its absence. The qualitative variable gender was characterized by “male” and “female”, race was characterized by “black” and “caucasian”, and in the variable smoking, the description “ex-smoker” was added, which included cases that quit smoking a minimum of 12 months before the exam. In the variable other cardioembolic heart diseases (sinus node diseases, valvulopathies, endocarditis, presence of prosthetic valves, cardiomyopathies, congenital heart diseases, and iatrogenic causes) the main potential embolic heart diseases except AF were included as “yes”. The data corresponding to the cervical Doppler ultrasound findings were grouped into carotid axes (includes the mean value of left and right axis results) and vertebral arteries (includes the mean value of left vertebral artery and right vertebral artery results) and were categorized into normal exams, mild stenosis (< 50%) and moderate stenosis (50% - 69%), significant stenosis (70% - 89%) and pre-occlusive stenosis (90% - 99%), occlusions, atheromatous changes without hemodynamic repercussions, occlusions and others (where tortuosity, hypoplasia, etc. were included). The quantitative variables age and IMI had their corresponding numerical values described, with the IMI presented in millimetres (mm) and classified as normal when the numerical value is ≤ 0.9 mm, thickening when found between 1 mm and 1.3 mm, and atheroma when the numerical value is ≥ 1.4 mm [13,14].

To quantify the degree and severity of carotid stenosis, the “North American Symptomatic Carotid Endarterectomy Trial” (NASCET) classification was used [15,16].

Ethnicity (“Caucasian” or “Black”) was not analysed, as there was only one black individual among 5884 Caucasian individuals.

Statistical analysis

The G*power Software (G*power 3.1, University of Düsseldorf, Düsseldorf, Germany) was used to analyse conditions with a sample of 5881 individuals: effect size $f = 0.25$, α error probability = 0.05 and power ($1 - \beta$ error probability) = 0.95.

An individual descriptive analysis of each variable was carried out. For continuous variables, mean and standard deviation were calculated; categorical variables were presented as a percentage. Testing the normality was performed using the Kolmogorov-Smirnov and Shapiro-Wilk tests.



For the evaluation of dependency relationships, the test according to the nature of the variables was used, including the Chi-square test; ETA test, and C-Pearson. p - values < 0.05 were considered statistically significant.

Statistical analysis was performed using SPSS statistical data analysis software (Statistical Package for the Social Sciences, IBM SPSS Statistics®23) in the Windows 10® version.

Ethical considerations

The study under analysis was carried out using a pre-existing database in an ultrasound laboratory at a center hospital, so there was no requirement in applying for the participants' consent, but it required the approval of the hospital's ethics committee (Act N°. 47).

To guarantee the confidentiality and anonymity of all data, they were coded sequentially and numerically, without any personal identification, respecting the ethical principles contained in the Declaration of Helsinki [17].

Results

A total of 5885 individuals were evaluated, of which 41.8% were female and 58.2% were male, with a mean age of 76.59 years \pm 6.686 and a maximum of 98 years.

Risk factors

With regard to modifiable risk factors, 81.3% of individuals had hypertension and 18.7% were normotensive. Dyslipidemia was present in 51.3% of individuals. With diabetes mellitus, 31.4% and most individuals did not report smoking habits (82.7%), 5.8% were smokers and 11.5% were ex-smokers. AF had a percentage of 16.6% and 3.6% with other cardioembolic heart diseases (Table 1).

Exam results

The minimum value found for the IMI was 0.5 mm and the maximum was 1.3 mm with an average of 0.962 mm (\pm 0.170). About 46.6% of the individuals had increased IMI.

In the evaluation of the carotid axes by Doppler ultrasound, 13.6% of the exams were normal, 64.2% had atheromatous plaques, without hemodynamic repercussions; 6.1% atheromatous plaques conditioning moderate stenosis, 5.4% atheromatous plaques conditioning significant pre-occlusive stenosis, and 3.3% with occlusion (by plaques, thrombus or atherothrombosis). The remainder (7.4%) had other alterations or variants of normality. Regarding the vertebral arteries, it was observed that most of the exams were normal (93.6%), 1.2% had atheromatous plaques, without hemodynamic repercussions, 0.3% atheromatous plaques with moderate stenosis, 0.3% atheromatous plaques conditioning significant stenosis to pre-occlusive, and 2.4% with occlusion. The remainder (2.2%) had other alterations or variants of normality (Table 2).

To analyse the relationship of dependence between the various risk factors referred to and alterations in the carotid

Table 1: Frequencies and percentages of risk factors ($n = 5885$).

Variable		Absolut frequencies	Percentage (%)
Family history of stroke	No	5816	98.8%
	Yes	69	1.2%
Family history of AMI	No	5844	99.3%
	Yes	41	0.7%
Hypertension	No	1098	18.7%
	Yes	4787	81.3%
Dyslipidemia	No	2867	48.7%
	Yes	3018	51.3%
Diabetes mellitus	No	4036	68.6%
	Yes	1849	31.4%
Smoke habits	No	4864	82.7%
	Yes/ex-smoker	1021	17.3%
AF	No	5025	85.4%
	Yes	860	14.6%
Other cardioembolic heart diseases	No	5675	96.4%
	Yes	210	3.6%

AMI: Acute Myocardial Infarction; AF: Atrial Fibrillation.

Table 2: Frequencies and percentages of cervical Doppler ultrasound results ($n = 5885$).

Variables	Absolut frequencies	Percentage (%)
Carotid axes		
Normal	803	13.6%
Atheromatous plaques, without hemodynamic repercussions	3776	64.2%
Moderate stenosis	359	6.1%
Pre-occlusive stenosis	315	5.4%
Occlusion	197	3.3%
Other	435	7.4%
Vertebral axes		
Normal	5507	93.6%
Atheromatous plaques, without hemodynamic repercussions	36	1.2%
Moderate stenosis	15	0.3%
Pre-occlusive stenosis	14	0.3%
Occlusion	139	2.4%
Other	174	2.2%

and vertebral axes, crosstabs, and statistical tests were carried out and percentages were compared, in which significant correlations were verified between genders and the presence of increased IMI and atheromatous alterations in the Doppler ultrasound. At the level of the carotid and vertebral axes ($p = 0.000$), there was a higher prevalence of IMI and alterations such as atheromatous plaques with or without stenosis in males compared to females. Regarding age, there is also a positive correlation with the IIM, which tends to have higher values with age. It should also be highlighted the presence of statistical significance between a family history of AMI and the presence of atheromatous disease in more advanced stages of atheromatous disease or other cervical arterial changes (Tables 3,4).

Regarding modifiable risk factors, hypertension, DM, and dyslipidemia stand out, which only showed statistical significance with the presence of alterations in the carotid



axes and vertebral arteries, with hypertensive, diabetic, or dyslipidemic patients being those who presented more changes. Smoking, on the other hand, showed a positive correlation with atheromatous disease in earlier stages (increased IMI) and in more advanced stages (presence of plaques with or without stenosis). AF had a significant correlation with the absence of atherosclerotic changes in the referred arteries (Tables 3,5).

Discussion

The aim was to analyse the presence of atheromatous disease in a group of individuals aged 65 years or over in this hospital sample and to understand its correlation with vascular risk factors. It is described in the literature that advanced age is one of the most important risk factors for the development of

Table 3: Crosstabs between non-modifiable risk factors and modifiable risk factors with cervical ecoDoppler results (n = 5885).

Variables			Cervical Doppler results		
			Normal (%)	Pathologic with hemodinamic repercurssion < 70%	Pathologic with hemodinamic repercurssion > 70%
Non-modifiable risk factors	Sex	female	6.9%	32.6%	2.3%
		male	5.8%	46.7%	5.7%
	Family history of stroke	No	12.6%	78.2%	8%
		Yes	0.2%	0.9%	0%
	Family history of AMI	No	12.6%	78.8%	7.9%
		Yes	0.1%	0.6%	0%
Hypertension	No	3.4%	13.9%	1.4%	
	Yes	9.3%	65.3%	6.7%	
Dyslipidemia	No	6.9%	37.9%	3.9%	
	Yes	5.7%	41.5%	4.1%	
Diabetes mellitus	No	9.5%	53.5%	5.6%	
	Yes	3.2%	25.8%	2.4%	
Smoke habits	No	11.3%	65.1%		
	Yes/ex-smoker	1.3%	14.2%	1.8%	
Modifiable risk factors	AF	No	11.1%	67.3%	7%
		Yes	1.6%	12%	1%
	Other cardioembolic heart diseases	No	12.4%	76.3%	7.7%
		Yes	0.3%	3%	0.3%

Legend: AMI – acute myocardial infarction; AF – atrial fibrillation.

Table 4: Relationship between non-modifiable risk factors and cervical Doppler ultrasound results (n = 5885).

Variable pairs		p-value
	sex vs. IMI	0
	sex vs. carotid axes/vertebral axes	0.000/0.000
	age vs. IMI*	0.002
	age vs. carotid axes/vertebral axes**	0.118/0.063
	family history of stroke vs. IMI	0.895
	family history of stroke vs. carotid axes/vertebral axes	0.083/1
	family history of AMI vs. IMI	0.928
Non-modifiable risk factos	family history of AMI vs. carotid axes/vertebral axes	0.343/0.021

AMI: Acute Myocardial Infarction; IMI: Intima-Media Index; p - value calculation: Chi-square; * C-Pearson test; ** ETA

Table 5: Relationship between modifiable risk factors and cervical Doppler ultrasound results (n = 5885).

Variable pairs		p - value
Modifiable risk factors	arterial hypertension vs. IMI	0.183
	arterial hypertension vs. carotid axes/vertebral axes	0.000/0.016
	DM vs. IMI	0.316
	DM vs. carotid axes/vertebral axes	0.000/0.039
	dyslipidemia vs. IMI	0.418
	dyslipidemia vs. carotid axes/vertebral axes	0.000/0.131
	smoke habits vs. IMI	0.000
	smoke habits vs. carotid axes/vertebral axes	0.000/0.000
	AF vs. IMI	0.103
	AF vs. carotid axes/vertebral axes	0.000/0.743
	other cardioembolic heart diseases vs. IMI	0.412
other cardioembolic heart diseases vs. carotid axes/vertebral axes	0.745/0.270	

AF: Atrial Fibrillation; DM: Diabetes Mellitus; IMI: Intima-Media Index; p-value calculation: Chi-square

atherosclerosis, constituting a public health problem because in the more advanced age group (> 65 years) there is greater frailty, number of diseases and risk factors for atherosclerotic disease.

In this sample, there was a higher prevalence of males, as described in the results of other studies related to atheromatous disease and risk factors [5,18]. According to some authors, atherosclerosis is the most frequent etiology and affects the coronary arteries, resulting in angina pectoris and/or acute myocardial infarction affects the arteries of the limbs resulting in peripheral vascular disease, and affects the extra and/or intracranial arteries, resulting in Cerebrovascular Accident (CVA) [6,7].

As for modifiable risk factors, it should be noted that most individuals are hypertensive (81.3%), about half of the sample have dyslipidemia (51.3%) and about 1/3 are diabetic. The average prevalence of hypertension found in the study is in line with what was expected and already reported in data from 2015, which refer to a prevalence of hypertension in the total Portuguese population of 36%, with a tendency to increase with advancing age [19]. Also, the result of dyslipidemia and diabetes are in line with what is described in the consulted literature. An epidemiological study by Mariano, et al. (2015), refers to a prevalence of 52% of dyslipidemia in the Portuguese population [20]. According to Correia, et al. (2009) in Portugal, more than a quarter of the population has diabetes mellitus [21]. According to the guidelines of the Sociedade Brasileira de Diabetes and the annual report of the national diabetes observatory in Portugal (2009), there is an increase in the prevalence of diabetes mellitus with the aging of individuals, thus concluding that the higher the average age, the greater the increase in the percentage of the disease [21].

As for other factors such as AF and smoking, its low prevalence may be related to a higher frequency of smoking in younger age groups and with better financial conditions [22] and AF, despite being more prevalent in the elderly, is found to be higher after the age of 80 [23], diverging from our sample, which has a relatively lower average age, as well as the lack of knowledge on the part of patients about some co-morbidities.

Regarding Doppler echo findings, less than a quarter (13.6%) of the sample showed normal carotid arteries and almost all (93.6%) had normal vertebral arteries. This agrees with the literature, since the presence of pathology is more frequent in larger calibre vessels that are subject to more turbulent flows that condition intimal lesions, increasing the predisposition for the development of the atherosclerotic process. In this exam, the measurement of the carotid wall IIM value is an important parameter in the cerebrovascular area, since it is a predictor of atherosclerotic disease [14]. Naqvi, et al. (2014) evaluated the intima-media index and atherosclerotic plaque in the assessment of cardiovascular risk and concluded that only the IIM value improves the predictive power of cardiovascular risk factors, but when the measurement is associated with the carotid bifurcation, it improves the prediction of disease atherosclerosis and stroke [24].

According to Blaha, et al. (2014), if the objective is to calculate the risk of a stroke, the ultrasound image with measurement of the IMI must be used [5]. In this sample, about half (46.6%) presented an increased IMI, that is, it presents signs of the atherosclerotic process in the initial phase. More than half of the sample (64.2%) has atheromatous disease in the carotid axes, but without hemodynamic repercussions in most individuals, which is in line with the fact that atherosclerosis is a chronic pathology with slow evolution that worsens with increasing age [25],

When analysing the relationship of dependence between the various risk factors referred to and changes in the carotid and vertebral axes, a significant relationship was observed between increased IMI and the presence of alterations such as atheromatous plaques with or without stenosis in males and in individuals with older ages, corroborating with previously mentioned studies [5,17,25]. A study carried out in 2008 reports that the IMI is associated with ages over 65 years [25]. In a meta-analysis carried out in 2017, it was concluded that the measurement of the thickness of the intima and media layers of the carotid arteries for assessment of cardiovascular risk, the IIM, increases with increasing age and male sex, due to the fact that with increasing age there is a greater development of multiple risk factors in the same individual that will cause to the increase of the IIM in relation to the younger ones [9]. In 2018, in a study with 84,880 individuals, a higher prevalence of carotid atherosclerosis was observed in males compared to females and increased with age [26]. as well as in a previous study carried out in 1999 [17]. Another study that evaluated the predictors of IMI and progression of atherosclerotic plaque in the Chinese population, found an association between increased IMI and progression of atherosclerotic plaque with gender, but a higher prevalence of these alterations was identified in females [27]. This and other studies point to a possible variability related to ethnicity and the region where the studies were carried out, since socioculturally there are customs where men play roles and have different habits (food, alcohol, sedentary lifestyle, education, profession, and information on health) [27,28]. In this sample the number of non-caucasian individuals was very small, thus these results are in agreement with the studies consulted with samples of similar characteristics.

Regarding modifiable risk factors, hypertensive, diabetic, or dyslipidemic individuals showed statistical significance with the presence of changes in the carotid axes and vertebral arteries (more advanced stages of the disease). The findings of this study are in line with a study carried out in 2010, where the development of atheroma plaques in diabetic and hypertensive patients was evaluated, which showed that associations between the formation of atheroma plaques and arterial hypertension are observed in the literature [29]. In a study Chinese epidemiological study carried out in 2018, individuals were evaluated ultrasonographically for carotid atherosclerosis, showing a significantly high rate of hypertension [26]. The more advanced age groups are more prone to the presence of modifiable risk factors such as hypertension, which constitutes one of the main risk factors for atherosclerotic disease, and it is expected that its presence, particularly in more advanced age groups, will influence the presence of atheromatous disease in medium and large calibre vessels, namely in areas of bifurcations or tortuosities that are very likely to damage the artery arterial wall and form atheromatous plaques [30]. Some studies have shown There are somewhat divergent results with regard to the relationship between hypertension and the different stages of the disease. In a study based on the evaluation of carotid atherosclerosis by echo-Doppler and the association with cardiovascular risk factors, arterial hypertension was not associated with carotid atherosclerosis, but with carotid intima-media thickening [25]. However, given the lower value (63 years) of the mean age of the sample in the study carried out may justify the absence of a positive relationship with more advanced stages of the disease and only being positive with the earlier stage.

As for dyslipidemia and diabetes, studies show that individuals with dyslipidemia have a higher prevalence of carotid atherosclerosis [28]. A meta-analysis revealed that they found 10 articles in which patients with dyslipidemia have a higher risk of developing carotid plaque and seven articles point to the fact of diabetic patients have a higher prevalence of atherosclerotic plaque than non-diabetic patients [30]. As previously mentioned, just like hypertension, diabetes mellitus is associated with atherosclerotic plaques. However, another study did not find statistical evidence of the association between carotid atherosclerosis and diabetes mellitus [25]. The literature presents inconsistent results on the subject [29]. The disagreement between studies is justifiable since it is described that diabetes mellitus is a risk factor for the development of atherosclerosis, mainly atherosclerosis of the microcirculation, thus less frequent direct involvement with large-calibre arteries and arterial bifurcations [31,32].

Smoking in this sample showed a positive correlation with atheromatous disease in the earlier stages of the disease (increased IMI) and in the more advanced stages (presence of plaques with or without stenosis). This corroborates the studies, where smoking is associated with carotid alterations [25,26].

Unlike in this sample, AF had a significant correlation with normal exams. This is in line with some research studies



previously carried out. AF and other cardioembolic diseases are the main cause of stroke, most of the time not due to disease of the local arterial wall, but due to the formation of thrombi of cardiac origin [33].

The main limitation of this study is the retrospective inclusion of individuals who received cervical Doppler, excluding all the patients that did not do it, which may introduce bias in the statistical results.

Conclusion

There are no published data on our Alentejo population, a very old one with particular diet habits. In this study, it was possible to characterize the correlation of non-modifiable risk factors such as advanced age and male sex with the increased occurrence of IMI. Male gender and modifiable risk factors such as hypertension, diabetes, dyslipidemia, and smoking are related to more advanced stages of atherosclerotic disease, with the presence of atheromatous plaques with and without hemodynamic repercussions.

References

- Cole JW. Large Artery Atherosclerotic Occlusive Disease. *Continuum (Minneapolis, Minn)*. 2017 Feb;23(1, Cerebrovascular Disease):133-157. doi: 10.1212/CON.0000000000000436. PMID: 28157748; PMCID: PMC5898962.
- HOLMAN RL, McGILL HC Jr, STRONG JP, GEER JC. The natural history of atherosclerosis: the early aortic lesions as seen in New Orleans in the middle of the 20th century. *Am J Pathol*. 1958 Mar-Apr;34(2):209-35. PMID: 13520905; PMCID: PMC1934740.
- Marques da Silva P, Lima MJ, Neves PM, Espiga de Macedo M. Prevalence of cardiovascular risk factors and other comorbidities in patients with hypertension in Portuguese primary health care populations: The PRECISE study. *Rev Port Cardiol (Engl Ed)*. 2019 Jun;38(6):427-437. English, Portuguese. doi: 10.1016/j.repc.2018.09.011. Epub 2019 Jul 12. PMID: 31307727.
- Polonia J, Martins L, Pinto F, Nazare J. Prevalence, awareness, treatment and control of hypertension and salt intake in Portugal: changes over a decade. The PHYSA study. *J Hypertens*. 2014 Jun;32(6):1211-21. doi: 10.1097/HJH.000000000000162. PMID: 24675681.
- Blaha MJ. The future of CV risk prediction: multisite imaging to predict multiple outcomes. *JACC Cardiovasc Imaging*. 2014 Oct;7(10):1054-6. doi: 10.1016/j.jcmg.2014.06.016. PMID: 25323166.
- Natural history of aortic and coronary atherosclerotic lesions in youth. Findings from the PDAY Study. *Pathobiological Determinants of Atherosclerosis in Youth (PDAY) Research Group. Arterioscler Thromb*. 1993 Sep;13(9):1291-8. doi: 10.1161/01.atv.13.9.1291. PMID: 8364013.
- Santos MG, Pegoraro M, Sandrini F, Macuco EC. Risk factors for the development of atherosclerosis in childhood and adolescence. *Arq Bras Cardiol*. 2008 Apr;90(4):276-283. English, Portuguese. doi: 10.1590/s0066-782x2008000400012. PMID: 18516390.
- Kitamura A, Iso H, Imano H, Ohira T, Okada T, Sato S, Kiyama M, Tanigawa T, Yamagishi K, Shimamoto T. Carotid intima-media thickness and plaque characteristics as a risk factor for stroke in Japanese elderly men. *Stroke*. 2004 Dec;35(12):2788-94. doi: 10.1161/01.STR.0000147723.52033.9e. Epub 2004 Nov 4. PMID: 15528460.
- Torres FS, Moreira CM, Vianna FF, Gus M. Intima-media thickness measurement for cardiovascular risk assessment. *Rev. bras. Hypertension*. 2007; 14(3) : 171-175.
- O'Leary DH, Bots ML. Imaging of atherosclerosis: carotid intima-media thickness. *Eur Heart J*. 2010 Jul;31(14):1682-9. doi: 10.1093/eurheartj/ehq185. Epub 2010 Jun 11. PMID: 20542989.
- Sousa-Uva M, Dias CM. Prevalence of stroke in the Portuguese population: data from the ECOS 2013/2014 sample.
- Perdigão C, Rocha E, Duarte JS, Santos A, Macedo A. Prevalence and distribution of the main cardiovascular risk factors in Portugal—the AMALIA study. *Rev Port Cardiol*. 2011 Apr;30(4):393-432. English, Portuguese. PMID: 21815523.
- Casella IB, Presti C, Porta RM, Sabbag CR, Bosch MA, Yamazaki Y. A practical protocol to measure common carotid artery intima-media thickness. *Clinics (Sao Paulo)*. 2008 Aug;63(4):515-20. doi: 10.1590/s1807-59322008000400017. PMID: 18719764; PMCID: PMC2664129.
- Pignoli P, Tremoli E, Poli A, Oreste P, Paoletti R. Intimal plus medial thickness of the arterial wall: a direct measurement with ultrasound imaging. *Circulation*. 1986 Dec;74(6):1399-406. doi: 10.1161/01.cir.74.6.1399. PMID: 3536154.
- Ferguson GG, Eliasziw M, Barr HW, Clagett GP, Barnes RW, Wallace MC, Taylor DW, Haynes RB, Finan JW, Hachinski VC, Barnett HJ. The North American Symptomatic Carotid Endarterectomy Trial : surgical results in 1415 patients. *Stroke*. 1999 Sep;30(9):1751-8. doi: 10.1161/01.str.30.9.1751. PMID: 10471419.
- Ferguson GG, Eliasziw M, Barr HW, Clagett GP, Barnes RW, Wallace MC, Taylor DW, Haynes RB, Finan JW, Hachinski VC, Barnett HJ. The North American Symptomatic Carotid Endarterectomy Trial : surgical results in 1415 patients. *Stroke*. 1999 Sep;30(9):1751-8. doi: 10.1161/01.str.30.9.1751. PMID: 10471419.
- Diniz D, Correa M. The Helsinki Declaration: relativism and vulnerability. *Public Health Notebooks/Ministry of Health, Oswaldo Cruz Foundation, National School of Public Health*. 2001; 17(3): 679-688. https://doi.org/10.1590/s0102-311x2001000300022.
- Joakimsen O, Bonna KH, Stensland-Bugge E, Jacobsen BK. Age and sex differences in the distribution and ultrasound morphology of carotid atherosclerosis: the Tromsø Study. *Arterioscler Thromb Vasc Biol*. 1999 Dec;19(12):3007-13. doi: 10.1161/01.atv.19.12.3007. PMID: 10591682.
- Rodrigues AP, Kislaya I, Graff-iversen S, Silva AC, Barreto M, Gil AP. Hypertension prevalence in Portugal: results from the First National Survey with Physical Examination (INSEF 2015)/Hypertension prevalence in Portugal: results from the first Portuguese Health Examination Survey 2015. 2017; (2): 11–4. ISSN: 0874-2928. www.insa.pt.
- Mariano C, Antunes M, Rato Q, Bourbon M. e_LIPID: characterization of the lipid profile of the Portuguese population. *National Institute of Health Doctor Ricardo Jorge, IP - Epidemiological Bulletin Observations*. 2015; 4(14): 7–10. ISSN: 2182- 88730874-2928.
- Correia LG, Boavida JM, Cardoso M, Duarte JS, Duarte R, Nunes JS. Diabetes: Facts and Figures 2016, 2017, and 2018 - Annual Report of the National Diabetes Observatory. Portugal. Portuguese Society of Diabetology. 2019.
- SNS. Tobacco consumption in Portugal. National Health Service. 2017. https://www.sns.gov.pt/noticias/2017/11/17/consumo-de-tabaco-em-Portugal/
- Bonhorst D. A new look at the prevalence of atrial fibrillation in Portugal: The Safira study. *Rev Port Cardiol (Engl Ed)*. 2018 Apr;37(4):315-317. English, Portuguese. doi: 10.1016/j.repc.2018.02.002. Epub 2018 Apr 22. PMID: 29685847.
- Naqvi TZ, Lee MS. Carotid intima-media thickness and plaque in cardiovascular risk assessment. *JACC Cardiovasc Imaging*. 2014 Oct;7(10):1025-38. doi: 10.1016/j.jcmg.2013.11.014. Epub 2014 Jul 16. PMID: 25051948.
- Freitas P, Piccinato CE, Martins WDP, Mauad Filho F. Carotid atherosclerosis evaluated by Doppler ultrasound: Association with risk factors and systemic



arterial disease. *J Vasc Bras.* 2008; 7(4): 298–307. <http://dx.doi.org/10.1590/S1677-54492009005000001>.

26. Wang X, Li W, Song F, Wang L, Fu Q, Cao S, Gan Y, Zhang W, Yue W, Yan F, Shi W, Wang X, Zhang H, Zhang H, Wang Z, Lu Z. Carotid Atherosclerosis Detected by Ultrasonography: A National Cross-Sectional Study. *J Am Heart Assoc.* 2018 Apr 5;7(8):e008701. doi: 10.1161/JAHA.118.008701. PMID: 29622590; PMCID: PMC6015437.
27. Huang LC, Lin RT, Chen CF, Chen CH, Juo SH, Lin HF. Predictors of Carotid Intima-Media Thickness and Plaque Progression in a Chinese Population. *J Atheroscler Thromb.* 2016 Aug 1;23(8):940-9. doi: 10.5551/jat.32177. Epub 2016 Feb 18. PMID: 26887219; PMCID: PMC7399290.
28. Zhang L, Fan F, Qi L, Jia J, Yang Y, Li J, Zhang Y. The association between carotid intima-media thickness and new-onset hypertension in a Chinese community-based population. *BMC Cardiovasc Disord.* 2019 Nov 27;19(1):269. doi: 10.1186/s12872-019-1266-1. PMID: 31775639; PMCID: PMC6882043.
29. Carvalho ACA de, Oliveira LS de AF, Melo DP de, Crusóé-Rebello I, Campos PSF. Development of atheromatous plaques in diabetic and hypertensive patients. *Rev Medical and Biological Sciences.* 2010; 9(1): 73. DOI: <https://doi.org/10.9771/cmbio.v9i1.4736>.
30. Ji X, Leng XY, Dong Y, Ma YH, Xu W, Cao XP, Hou XH, Dong Q, Tan L, Yu JT. Modifiable risk factors for carotid atherosclerosis: a meta-analysis and systematic review. *Ann Transl Med.* 2019 Nov;7(22):632. doi: 10.21037/atm.2019.10.115. PMID: 31930033; PMCID: PMC6944535.
31. Triches C, Schaan BDA, Gross JL, Azevedo MJ. Macrovascular complications of diabetes mellitus: Clinical, diagnostic and management peculiarities. *Arq Bras Endocrinol Metabol.* 2009; 53(6): 698–708. <https://doi.org/10.1590/S0004-27302009000600002>.
32. Amarante RDM, Castro R, Lage AV, Cisternas JR. Diabetes Mellitus as a risk factor in atherogenesis. *Medical Archives of Hospitals and Faculty of Medical Sciences of Santa Casa de São Paulo.* 2007; 52(3): 87–93.
33. Kamel H, Healey JS. Cardioembolic Stroke. *Circ Res.* 2017 Feb 3;120(3):514-526. doi: 10.1161/CIRCRESAHA.116.308407. PMID: 28154101; PMCID: PMC5312810.

Discover a bigger Impact and Visibility of your article publication with Peertechz Publications

Highlights

- ❖ Signatory publisher of ORCID
- ❖ Signatory Publisher of DORA (San Francisco Declaration on Research Assessment)
- ❖ Articles archived in worlds' renowned service providers such as Portico, CNKI, AGRIS, TDNet, Base (Bielefeld University Library), CrossRef, Scilit, J-Gate etc.
- ❖ Journals indexed in ICMJE, SHERPA/ROMEO, Google Scholar etc.
- ❖ OAI-PMH (Open Archives Initiative Protocol for Metadata Harvesting)
- ❖ Dedicated Editorial Board for every journal
- ❖ Accurate and rapid peer-review process
- ❖ Increased citations of published articles through promotions
- ❖ Reduced timeline for article publication

Submit your articles and experience a new surge in publication services (<https://www.peertechz.com/submission>).

Peertechz journals wishes everlasting success in your every endeavours.